

Supplemental Material to:

**Tanmaya Joshi, Malay Patra, Leone Spiccia
and Gilles Gasser**

**Di-heterometalation of thiol-functionalized peptide
nucleic acids**

**Artificial DNA: PNA & XNA 2013; 4(1)
<http://dx.doi.org/10.4161/adna.24019>**

**[http://www.landesbioscience.com/journals/artificialdna/
article/24019](http://www.landesbioscience.com/journals/artificialdna/article/24019)**

Supplementary Information:

Di-Heterometalation of Thiol-Functionalized Peptide Nucleic Acids

Tanmaya Joshi,^a Malay Patra,^a Leone Spiccia^b and Gilles Gasser^{a,}*

^{a)} Institute of Inorganic Chemistry, University of Zurich, Winterthurerstrasse 190, CH-8057 Zurich, Switzerland.

^{b)} ARC Centre of Excellence for Electromaterials Science and School of Chemistry, Monash University, Clayton, Victoria 3800, Australia

* Corresponding author: Email: gilles.gasser@aci.uzh.ch; Fax: +41 44 635 6803; Tel: +41 44 635 4630; WWW: www.gassergroup.com.




Figure S1. ¹H NMR spectrum of **C2** in CDCl_3 .




Figure S2. ¹³C NMR spectrum of **C2** in CDCl_3 .




Figure S3. ¹H NMR spectrum of **C3** in CDCl_3 .




Figure S4. ¹³C NMR spectrum of **C3** in CDCl_3 .




Figure S5. MALDI-TOF spectrum of **PNA1**.




Figure S6. LC-MS spectrum of **PNA1**.




Figure S7. LC-MS spectrum of PNA2.





Figure S8. LC-MS spectrum of PNA3. Note that the two major peaks in the MS spectrum have a similar MS pattern.




Figure S9. LC-MS spectrum of PNA4.




Figure S10. LC-MS spectrum of PNA5.





Figure S11. LC-MS spectrum of **PNA6 (NLS-PNA-lys)**.




Figure S12. IR spectrum of PNA3.




Figure S13. IR spectrum of PNA5.